



**Eur päisches  
Patentamt**

**European  
Patent Office**

**Office eur péen  
des brevets**

**Bescheinigung**

**Certificate**

**Attestation**

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

**BEST AVAILABLE COPY**

**Patentanmeldung Nr. Patent application No. Demande de brevet n°**

00128660.8

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**R C van Dijk**

DEN HAAG, DEN  
THE HAGUE, 10/12/01  
LA HAYE, LE



**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.:  
Demande n°: 00128660.8

Anmeldetag:  
Date of filing:  
Date de dépôt: 28/12/00

Anmelder:  
Applicant(s):  
Demandeur(s):  
SHAN-Beteiligungsgesellschaft m.b.H.  
8700 Leoben  
AUSTRIA

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:  
Allergy vaccines containing hybrid polypeptides

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

/

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE/TR  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

**BEST AVAILABLE COPY**

EPO - Munich  
52

**28. Dez. 2000**

December 28, 2000

SHAN-Beteiligungsgesellschaft m.b.H.  
Dirnböckweg 7/10  
A-8700 Leoben  
Österreich

---

**Allergy vaccines containing hybrid polypeptides**

---

Type I allergy is a genetically determined hypersensitivity disease affecting almost 500 million individuals worldwide. Immediate symptoms (allergic rhinoconjunctivitis, asthma, anaphylactic shock) as well as late symptoms (atopic dermatitis, certain forms of allergic asthma bronchiale) are based on the recognition of allergens by IgE antibodies. Immediate symptoms result from the allergen-induced crosslinking of effector cell-bound IgE and the subsequent release of biological mediators (e. g., histamine, leukotrienes) whereas late symptoms can be caused by IgE-mediated presentation of allergens to T cells and eosinophil activation.

The only curative therapy approach, allergen-specific immunotherapy, is based on the systemic administration of allergens to patients in order to induce allergen-specific "unresponsiveness" (Noon, Lancet 1911, 1: 1572-1573; Bousquet et al. (1998) J Allergy Clin Immunol 102: 558-562; Durham and Till (1998) J. Allergy Clin. Immunol. 102, 157-164). Using conventional technologies it has not been possible to produce pure allergens for specific immunotherapy and hence, vaccination is performed with allergen extracts. These extracts consist of allergens and non-allergenic components which are difficult to

BEST AVAILABLE COPY

## BEST AVAILABLE COPY

standardize. Therefore patients cannot be treated according to their specific sensitization profile and anaphylactic side effects, due to the administration of allergenic material are frequently observed (Møllerup et al. (2000). Clin. Exp. Allergy 30, 1423-1429).

It is an objective of the present invention to provide an advantageous composition for the treatment or prevention of allergic disorders.

Surprisingly the inventors found that it is possible to generate hybrid allergens assembling the epitopes of immunologically distinct allergens for diagnosis and therapy of Type I allergy. The invention therefore relates to a hybrid polypeptide comprising at least two different allergenic proteins or fragments thereof wherein each fragment consists of at least eight consecutive amino acids of the respective allergenic protein.

The allergenic proteins from which the hybrid polypeptide is derived may be a plant allergen. The different allergenic proteins may be derived from a common source such as the pollens of a certain plant species. Several different allergenic proteins contained in the allergen source may be identified and used for the preparation of the hybrid polypeptide of the invention. The present invention, however, encompasses also a hybrid polypeptide comprising sequences derived from allergenic proteins from different sources. The term "allergenic proteins or fragments thereof" comprises also modifications of the allergens wherein the sequence of the naturally occurring allergen has been slightly modified by substitutions of single amino acids or nucleotides whereby the allergenic potential has been substantially maintained.

Allergen sources from which the allergenic proteins are derived may be major grass pollen, mite, bee venom or animal hair dander allergens. Specific examples of allergenic proteins are the group 1, group 2, group 4, group 5 group 6, group 11, group 12 and group 13 allergens of major grass pollen, Der p 1 and Der p 2 (mite), phospholipase from bee venom and Fel d 1 (cat).

In one embodiment the hybrid polypeptide comprises at least one complete allergenic protein. It may also comprise two different complete allergenic proteins. It is not only possible to combine different groups of allergenic grass pollen but also to combine allergenic proteins derived from different sources. In a particular embodiment all sequences

## BEST AVAILABLE COPY

of the hybrid polypeptide derived from allergenic proteins represent complete allergenic proteins.

In another embodiment the hybrid polypeptide comprises at least one fragment of an allergenic protein wherein the fragment consists of at least eight consecutive amino acids of the respective allergenic protein. Preferably the fragment consists of at least 12, more preferably of at least 20 and most preferably of at least 30 consecutive amino acids of the respective allergenic proteins. In another embodiment all amino acid sequences derived from allergenic proteins are fragments of at least eight consecutive amino acids of the respective allergenic proteins from which they are derived. The preferred length of these fragments is at least 12, more preferably at least 20, most preferably at least 30 consecutive amino acids of the respective allergenic protein.

When fragments of allergenic proteins are employed it is possible to prepare a hybrid polypeptide comprising only fragments which have an allergenic activity which is lower compared with the respective allergenic proteins from which they are derived. This effect may be due to the destruction of epitopes by modified secondary or tertiary structure of the fragment compared with the full length protein. In the most preferred embodiment, the hybrid polypeptide has an allergic activity which is lower than the allergenic activity of each of the allergenic proteins from which the hybrid polypeptide is derived. Usually, the allergenic activity of the hybrid polypeptide is less than 50% of that of each of the allergenic proteins from which the hybrid polypeptide is derived. In a particular embodiment, the hybrid polypeptide has substantially no allergenic activity.

According to the invention the allergenic activity of a sample is determined by determining the IgE antibodies which are induced in a test animal upon application of the sample. The allergenic activity is preferably defined in suitable in vitro or in vivo tests. A preferred in vitro test is the basophil histamine release assay as described in Vrtala et al., J. Clin. Invest. 1997, 99, pp. 1673-1681. Alternatively the allergenic activity is determined in a skin test as described in van Hage-Hamsten et al. J. Allergy Clin. Immunol. 1999, 104, pp. 969-977 or in Pauli et al. Clin. Exp. Allergy 2000, 30, pp. 1076-1084.

In one embodiment the hybrid polypeptide comprises two portions which are derived from two different allergenic proteins. The hybrid polypeptide of the invention, however, may

comprise three, four, five or even more portions each of which is derived from a different allergenic protein.

The hybrid polypeptide of the invention does not necessarily consist only of amino acid sequences derived from allergenic proteins. It is possible that artificial sequences (e.g. spacer sequences) are inserted between the units representing sequences from different allergenic proteins. It is also possible that the amino acid sequences of the naturally occurring allergenic proteins are modified, e.g. by genetic engineering to introduce mutations which reduce the allergenic activity of the fragment. It is also preferred that the hybrid polypeptide comprises a "tag" sequence which facilitates the purification of the hybrid polypeptide upon expression in a host cell. An example of a "tag" is the hexahistidine tag which allows purification by  $\text{Ni}^{2+}$  chelate chromatography. Other tags are known in the art.

The hybrid polypeptide of the invention may be prepared by several methods. In one embodiment the polypeptide is prepared by expressing a polynucleotide in a host cell. The host cell may be a procaryotic or a eucaryotic cell. If procaryotic cells are used the host is preferably E.coli. Examples of eucaryotic cells are yeast, insect cells or cell lines like CHO cells. After introducing a suitable polynucleotide encoding the polypeptide of the invention into a host cell the host cell is cultured under conditions such that the polypeptide is expressed in the cell. The polypeptide may be secreted by the cell or accumulate inside the cell. Known purification methods can be used to recover the hybrid polypeptide from the cell or from the culture medium.

The invention also encompasses the preparation of the hybrid polypeptide by chemical synthesis such as solid phase synthesis.

The invention further concerns a polynucleotide encoding a hybrid polypeptide according to the invention. Due to the degeneracy of the genetic code many different polynucleotide molecules may encode a single polypeptide. The polynucleotide of the invention preferably is an expression construct for obtaining the polypeptide after expression in host cells. The expression construct may further comprise components which are generally known in the art such as promoter sequences, genes encoding resistance factors against antibiotics, a replication origin etc.

## BEST AVAILABLE COPY

The invention further relates to a cell transfected or transformed with a polynucleotide of the invention. The cell may be a eucaryotic cell or a procaryotic cell. Eucaryotic cells may be transfected by a method known per se such as calcium phosphate mediated transfection, electroporation, lipofection etc.

The invention further relates to a pharmaceutical composition containing a polypeptide, polynucleotide or a cell according to the invention. The pharmaceutical composition may further contain a pharmaceutically acceptable carrier or diluent such as a buffer or salt solution. Preferably the pharmaceutical composition of the invention is a vaccine composition. In a particular embodiment the pharmaceutical composition further contains an adjuvant such as  $Al(OH)_3$ .

The invention also relates to a method for the preparation of a hybrid polypeptide of the invention. The method comprises providing a polynucleotide encoding a hybrid polypeptide, introducing said polynucleotide into a host cell, culturing the host cell thus obtained under conditions such that the hybrid polypeptide is expressed, and recovering the expression product from the cell. The polynucleotide may be prepared by methods known in the art, it is preferred that PCR technology is used to prepare the polynucleotide encoding the hybrid polypeptide.

The invention further concerns the use of a hybrid polypeptide, of a polynucleotide or of a cell of the invention for the preparation of a medicament for the treatment of an allergic disorder.

Such a medicament may be composed of the polynucleotide encoding a hybride vaccine which can be used directly for DNA-based vaccination against Type I allergy. The recombinant or synthetic hybrid polypeptide may be used to prepare formulations for the oral, sublingual or parenteral treatment of Type I allergic disorders as they are now routinely used for immunotherapy. Examples are formulations for sublingual immunotherapy or adjuvant bound hybrid polypeptides for injection immunotherapy. Possible applications comprise also cell based forms of immunotherapy which may be based on e.g., dendritic cells or other antigen presenting cells. Those cells are transformed and express the antigen in vivo. Preferably autologous cells transformed with suitable vectors are used.

One mode of applications may be the subcutaneous injection of adjuvant-bound hybrid polypeptides. Another possibility is the oral or nasal administration of the hybrid polypeptide in order to induce immunological tolerance or anergy against the components of the hybrid polypeptides. All these possible formulations can be prepared according to rules (dosage, adjuvants, schemes of administration) which are known to the person skilled in the art.

The invention also relates to the use of a hybrid polypeptide, of a polynucleotide, or of a cell of the invention for the preparation of a medicament for prophylactic vaccination or tolerance induction. Prophylactic administration of hybrid polypeptides means the administration of the polypeptide to individuals, preferentially children, who do not yet suffer from Type I allergy in order to induce a state of immunological tolerance, anergy or non-responsiveness, or a protective immunity against the components of the hybrid vaccine. This may be achieved by the various protocols outlined for treatment of an established allergic disorder. The prophylactic treatment may be performed with hybrid polypeptides consisting of hybrid polypeptides or the polynucleotides coding for the hybrid polypeptides as outlined before.

In a further embodiment the invention relates to the use of a hybrid polypeptide of the invention for the detection of antibodies against an allergenic protein in a sample. The antibody may be an IgM, IgE, IgG or IgA antibody. The concentration of the antibody may be determined from a sample which has been obtained from a body fluid. The sample may be derived from animals or humans. Such tests may use a solid phase-immobilized hybrid polypeptide or the hybrid polypeptide in the fluid phase. Examples for such tests may be ELISA tests. Westernblotting tests or any other test where the hybrid polypeptide is immobilized to bind specific antibodies out from the test sample. Alternatively the hybrid polypeptide is added directly to the antibody-containing fluid in order to adsorb specific antibodies as e.g., is done for competitive immunological assays.

The polypeptide of the invention may also be used for cellular tests such as a T-cell proliferation test, mediator release test etc. The hybrid polypeptide may be exposed to various types of cells in order to elicit measurable responses. Such responses may comprise the release of histamine or other mediators (e.g., leukotriens, serotonin, ECP) in the case of allergic effector cells (e.g., basophils mast cells, eosinophils). In another type of



assay the proliferation or death (e.g., apoptosis) of cells may be measured e.g., by the uptake of  $^3\text{H}$  Thymidine or any other suitable assay. Such cells may be T cells. Furthermore, hybrid polypeptides may be used to induce the release of cytokines or other immunologically relevant substances (e.g., from T cells) that can be measured. Furthermore, they may be used for antigen presentation assays. The hybrid polypeptide of the invention may also be used for diagnostic screening purposes. Hybrid polypeptides may be also used for in vivo provocation testing. Such tests may comprise skin testing (e.g., skin prick or intradermal testing), nasal provocation testing, all forms of food challenge testing or bronchial provocation testing.

Since hybrid polypeptides can contain epitopes of unrelated allergens they may be used for diagnostic screening tests (in vitro, in vivo as outlined above) in order to detect sensitization or unresponsiveness against one of the components of the hybrid polypeptide. This may allow to provide the physician with a diagnostic test which is suited to screen for sensitized patients in a fast way. Currently such tests (e.g., Phadiatop, Pharmacia, Uppsala, Sweden) consist of a mixture of non-covalently bound allergens which are coupled to one carrier. Therefore, the coupling rate of the individual components is difficult to control whereas a hybrid polypeptide would represent a preferential formulation for the preparation of such a screening test.

More than 40% of allergic patients are sensitized to grass pollens from different species. Grasses are also important allergen sources because of their worldwide distribution and heavy pollen production. They contain a variety of different allergenic components which occur as crossreactive allergens in different monocots. The inventors demonstrated that it is possible to generate by recombinant DNA technology hybrid allergens which consist of immunologically unrelated allergens and their epitopes.

In a first step frequent co-sensitizations were determined using purified recombinant timothy grass pollen allergens to select the most frequently recognized allergen combinations for the hybrid allergen approach. According to the pilot experiments hybrids consisting of immunologically unrelated major timothy grass pollen allergens were engineered. They comprised hybrids of the major timothy grass pollen allergens Phl p 5/Phl p 1, Phl p 2/Phl p 6 and a combination of all four molecules. As demonstrated for Phl p 2/Phl p 6 we found no evidence that a certain order of engineering the individual components would affect the

presentation of epitopes. Both, the rPhl p 2/Phl p 6 hybrid and the rPhl p 6/Phl p 2 hybrid preserved the relevant B and T cell epitopes of the components. The giant consisting of all 4 allergens also contained the relevant epitopes of the single components.

The hybrid allergen approach will open new avenues for diagnosis and treatment of Type I allergy. Hybrids consisting of the most frequently recognized allergens with preserved epitopes represent candidate molecules for in vitro as well as in vivo allergy screening tests. The giant described in the present application will likely identify more than 95% of timothy grass pollen allergic patients when used for diagnosis, because it resembles most of the relevant IgE epitopes present in natural timothy grass pollen extract. Molecules such as the giant can be produced also for other allergen systems (e. g., mite allergy). The advantage of using hybrid allergens for diagnosis is that they can be produced in a well controlled manner. In contrast to natural allergen extracts, which consist of a difficult to standardize mixture of allergens and non-allergenic materials, hybrid allergens will contain the relevant epitopes in a well defined ratio.

Recombinant hybrid allergens can also be used for immunotherapy. They can either be composed of the biologically active components or of hypoallergenic allergen units. Hybrid allergens covering frequent co-sensitization patterns may then be used for the treatment of allergies to complex allergen sources, e.g., grass pollen allergy because they resemble the relevant epitopes of the complete extract. Since the hybrid allergens consist of defined components it will be possible to produce well defined formulations for vaccination treatment. Hybrid allergens produced out of hypoallergenic variants will be tolerated in high doses and thus can be used like vaccines regularly used for prevention of viral infections. In this respect it may be considered to use hybrid allergens containing the epitopes of the most relevant allergens for prophylactic vaccination or tolerance induction even in not yet sensitized individuals.

Description of the tables and figures:

Table 1 shows the percentage inhibition of IgE binding to recombinant Phl p 2 and Phl p 6 achieved after preadsorption of sera from 20 patients with the hybrid allergens (rPhl p 2/rPhl p 6, rPhl p 6/rPhl p 2) or the recombinant allergens (rPhl p 2, rPhl p 6).

Table 2 shows the percentage inhibition of IgE binding to recombinant Phl p 1, Phl p 5, and Bet v 1 achieved after preadsorption of sera from 20 patients with the hybrid allergen rPhl p 5/rPhl p 1 or the recombinant allergens (rPhl p 1, rPhl p 5, and rBet v 1).

Figure 1 shows the frequency of co-sensitization to timothy grass pollen allergens rPhl p 1, rPhl p 2, rPhl p 5, and rPhl p 6 as determined in a population of 110 grass pollen allergic patients.

Figure 2 outlines the construction of hybrid vaccines. The fusion of the two genes (e.g. Phl p 5, Phl p 1) is mediated by a two-step PCR reaction, creating overlapping ends in the first PCR reaction followed by the amplification of both sequences in a second PCR step.

Figure 3 shows the result of example 3. Purified timothy grass pollen allergens (rPhl p 1, rPhl p 2, rPhl p 5, and rPhl p 6) and hybrids of them (rPhl p 5/rPhl p 1, rPhl p 2/rPhl p 6, and rPhl p 6/rPhl p 2) were loaded on a SDS polyacrylamide gel and stained with Coomassie brilliant blue. Molecular weights are displayed at the left margin.

In figure 4 (A) a purified recombinant hybrid consisting of Phl p 6/Phl p 2/Phl p 5/Phl p 1 was dotted on nitrocellulose filters. In row (B) rPhl p 2 (lane 1,3,4, and 5) and rPhl p 5 (lane 2) was immobilized. Filters were incubated with rabbit antisera against rPhl p 1 (lane 1), rPhl p 2 (lane 2), rPhl p 5 (lane 3), and rPhl p 6 (lane 4). The filter in lane 5 was exposed to a mouse monoclonal anti-His-tag antibody.

The following examples further illustrate the invention:

#### **Example 1: Determination of frequent co-sensitization patterns**

Grasspollen contain a variety of immunologically distinct and related allergens. These components are recognized at different frequencies and intensities. Recombinant timothy grass pollen allergens, rPhl p 1, rPhl p 2, rPhl p 5, and rPhl p 6, were previously identified as the most important components which resemble most of the IgE epitopes of natural grass pollen extracts (Niederberger et al. (1998). IgE antibodies to recombinant pollen allergens (Phl p 1, Phl p 2, Phl p 5, and Bet v 2) account for a high percentage of grass pollen-specific IgE. J Allergy Clin Immunol 101: 258-264). The results of the evaluation of

co-sensitizations in 110 grass pollen allergic patients to combinations of the recombinant timothy grass pollen allergens are displayed in Figure 1. ELISA testing with purified recombinant allergens showed that 60% percent of the patients were co-sensitized to rPhl p 1 and rPhl p 5, 30% contained IgE antibodies to rPhl p 2 and rPhl p 6 and 30% reacted simultaneously to all four recombinant allergens (Figure 1). Previous competition studies had indicated that a mixture of the above four allergens contained the majority of IgE epitopes present in 8 different monocot pollens (grasses, corn) (Laffer et al. (1996) Comparison of recombinant timothy grass pollen allergens with natural extract for diagnosis of grass pollen allergy in different populations. *J Allergy Clin Immunol*; 98: 652-658).

Patients sera, antisera and recombinant allergens:

Sera of 110 allergic patients were characterized by a positive timothy grass pollen RAST (radioallergosorbent test) result and by determination of IgE antibodies to timothy grass pollen extract by Western blotting as described (Niederberger et al. (1998) *J Allergy Clin Immunol* 101: 258-264). The recombinant timothy grass pollen allergens rPhl p 1, rPhl p 2, rPhl p 5, and rPhl p 6 were purified as described previously (Vrtala et al. (1995) *J. Allergy Clin. Immunol* 97: 781-787; Vrtala et al. (1999) *J. Immunol.* 163:5489-5496). The rabbit anti-rPhl p 1, anti-rPhl p 2, anti-rPhl p 5, and anti-rPhl p 6 antisera were raised against purified rPhl p 1, rPhl p 2, rPhl p 5, and rPhl p 6 using CFA (Charles River, Kisslegg, Germany).

## **Example 2: Construction of recombinant hybrid allergens**

The protocol for the construction of recombinant hybrid allergens is displayed in Figure 2. As exemplified for a hybrid consisting of rPhl p 5 linked to rPhl p 1, cDNAs coding for the components are amplified with suitable primer pairs (Figure 2: Phl p 5: w, x; Phl p 1: y, z) in order to create overlapping ends. In a subsequent second PCR reaction, a cDNA comprising both cDNAs is created with primers specific for the 5' end of the first cDNA (Phl p 5: w) and the 3' end of the second cDNA (Phl p 1: z) using both PCR products of the first reactions as templates. Using this technology we produced recombinant hybrids consisting of a combination of Phl p 5/Phl p 1, rPhl p 2/rPhl p 6, rPhl p 6/rPhl p 2 and of all four allergens (N-terminus-rPhl p 6-rPhl p 2-rPhl p 5-rPhl p 1-C-terminus), the latter referred to as the "giant".

Construction of expression plasmids for rPhl p 5/rPhl p 1-, rPhl p 2/rPhl p 6-, rPhl p 6/rPhl p 2-, and rPhl p 6/rPhl p 2/rPhl p 5/rPhl p 1 hybrid proteins:

The cDNAs of Phl p 5 and Phl p 1 were obtained by polymerase chain reaction using the primers 5' GGA ATT CAT ATG GCC GAT CTC GGT TAC 3' (SEQ ID NO: 1) and 5' CGG GGT ACC GAC TTT GTA GCC ACC AGT 3' (SEQ ID NO: 2) for Phl p 5 and 5' CGG GGT ACC ATG ATC CCC AAG GTT CCC 3' (SEQ ID NO: 3) and 5' CGG GAT CCT CAG TGG TGG TGG TGG TGG TGC TTG GAC TCG TAG CTG GT 3' (SEQ ID NO: 4) for Phl p 1. The signal peptide of Phl p 5 was replaced by a NdeI restriction site, containing the ATG start codon at the 5' end of the coding region. A KpnI restriction site was introduced at the 3' end of Phl p 5 replacing the stop-codon. The Phl p 1 sequence, lacking the signal peptide, started with a KpnI restriction site at the 5' end. At the 3' end a nucleotide stretch coding for a Hexahistidine-tag was introduced, followed by a stop codon and a BamHI restriction site. The PCR products were inserted as NdeI/KpnI/BamHI fragment into a pET17b expression vector.

The Phl p 2/Phl p 6 and Phl p 6/Phl p 2 sequences were constructed using PCR based 'gene soeing' (17). In a first PCR reaction the cDNAs of Phl p 2 and Phl p 6 were amplified using the primers 5' GGA ATT CAT ATG GTG CCG AAG GTG ACG 3' (SEQ ID NO: 5), and 5' CGT GGC CTT CCC CAT AAG CTT CTC TTC TGG CGC GTA GGT 3' (SEQ ID NO: 6) for Phl p 2 and 5' AAG CTT ATG GGG AAG GCC ACG ACC 3' (SEQ ID NO: 7) and 5' C GGG ATC CTA GTG GTG GTG GTG GTG GTG CGC GCC GGG CTT GAC AGC 3' (SEQ ID NO: 8) for Phl p 6 to create overlapping ends. In a subsequent PCR reaction the two agarose gel purified PCR products (using QIAGEN gel purification kit) were used as templates and amplified using the primers 5' GGA ATT CAT ATG GTG CCG AAG GTG ACG 3' (SEQ ID NO: 9) and 5' C GGG ATC CTA GTG GTG GTG GTG GTG GTG CGC GCC GGG CTT GAC AGC 3' (SEQ ID NO: 10). The signal peptide of Phl p 2 was replaced by a NdeI restriction site at the 5' end, the stop codon at the 3' end was removed. In the Phl p 6 sequence, also lacking the signal peptide, a Hexahistidine tag was introduced followed by a stop codon and a BamHI restriction site. The resulting Phl p 2/Phl p 6 construct was inserted as a NdeI/BamHI fragment into the pET17b expression vector.

The Phl p 6/Phl p 2 sequence was constructed using the same method with the PCR primers 5' GGA ATT CAT ATG GGG AAG GCC ACG ACC GAG 3' (SEQ ID NO: 11) and 5'

CAC CTT CGG CAC CAT AAG CTT CGC GCC GGG CTT GAC AGC 3' (SEQ ID NO: 12) for Phl p 6 and 5' AAG CTT ATG GTG CCG AAG GTG ACG 3' (SEQ ID NO: 13) and 5' C GGG ATC CTA GTG GTG GTG GTG GTG GTG CTC TTC TGG CGC GTA GGT 3' (SEQ ID NO: 14) for Phl p 2 in the first PCR reaction and the primers 5' GGA ATT CAT ATG GGG AAG GCC ACG ACC GAG 3' (SEQ ID NO: 15) and 5' C GGG ATC CTA GTG GTG GTG GTG GTG GTG CTC TTC TGG CGC GTA GGT 3' (SEQ ID NO: 16) for the second PCR reaction. The resulting Phl p 6/Phl p 2 construct was inserted in the pET17b vector as a NdeI/BamHI fragment.

For the construction of the Phl p 6/Phl p 2/Phl p 5/Phl p 1 hybrid ("giant") the Phl p 6/Phl p 2 construct was taken as a template in a PCR reaction using the primers 5' GGA ATT CAT ATG GGG AAG GCC ACG ACC GAG 3' (SEQ ID NO: 17) and 5' GGG ATT TCC ATA TGC TCT TCT GGC GCG TAG G 3' (SEQ ID NO: 18) replacing the 3' Hexahistidine tag and the stop codon by an NdeI restriction site. The sequence was cloned into the pET17b vector, containing the Phl p 5/Phl p 1 construct as a NdeI fragment. The right orientation of the insert was examined by restriction digest.

### Example 3: Expression and purification of hybrid allergens

Recombinant hybrid allergens with a C-terminal hexahistidine tag were expressed in *E. coli* BL21 (DE3) and purified by Nickel affinity chromatography. The rPhl p 2/rPhl p 6 and rPhl p 6/rPhl p 2 hybrid were purified from the soluble cytoplasmic fraction of *E. coli* extracts whereas the rPhl p 5/rPhl p 1 hybrid and the "giant" were solubilized from the insoluble inclusion body fraction of *E. coli* in urea. Figure 3 shows a Coomassie-stained SDS-PAGE containing three of the hybrids and the single recombinant allergens. The molecular mass determined by SDS-PAGE corresponded to the molecular weight calculated for the hybrids on the basis of their deduced amino acid sequence (rPhl p 5/rPhl p 1:60kDa ; rPhl p 2/rPhl p 6 and rPhl p 6/rPhl p 2:22kDa ; "giant": 82kDa; data not shown).

#### Experimental protocol:

All constructs were expressed in *E. coli* BL21 (DE3). Cells were grown in LB-medium containing 100mg/l ampicillin to an OD<sub>600</sub> of 0.8. The expression of the recombinant proteins

was induced by adding isopropyl- $\beta$ -thiogalactopyranoside (IPTG) to a final concentration of 0.5mM. After 4 hours at 37°C cells were harvested by centrifugation.

rPhl p 2/rPhl p 6 and rPhl p 6/rPhl p 2 were expressed as soluble proteins. Cells were resuspended in lysis buffer (50mM NaH<sub>2</sub>PO<sub>4</sub>, 300mM NaCl, 10mM imidazole, pH 8.0) and lysed with an ultraturrax (Polytron, Kinematica AG, Switzerland). The lysate was centrifuged at 10,000xg for 30min at 4°C to pellet the cellular debris. The supernatant was loaded on a Ni-NTA column (QIAGEN), the proteins were eluted with 250mM imidazole and dialyzed against water.

rPhl p 5/rPhl p 1 and rPhl p 6/rPhl p 2/rPhl p 5/rPhl p 1 were expressed in the inclusion body fraction. Cells were resuspended in 10mM Tris, 0.1% Triton, pH 7.4 and lysed by adding lysozyme to 1mg/ml. After centrifugation at 14 000xg for 20min at 4°C the pellet was washed 4x with 5mM Tris pH 8.0, 0.05M NaCl, 0.25% Desoxicholate, 0.25mM  $\beta$ -Mercaptoethanol and once with 10mM Tris, pH 8.0, 3% isopropanol. Inclusion bodies were solubilized with 8M urea, 100mM NaH<sub>2</sub>PO<sub>4</sub>, 10mM Tris, pH 8.0. After centrifugation at 14 000xg for 20min the supernatant was loaded on a Ni-NTA column (Qiagen). The proteins were renatured on the column using a linear 6M-1M urea gradient in 500mM NaCl, 20% glycerol, 20mM Tris, pH 7.4) over a period of 1.5h (18). Proteins were eluted by addition of 250mM imidazole and dialyzed against PBS pH 7.4.

#### **Example 4: Hybrid allergens contain the relevant epitopes of their components**

Recombinant hybrid allergens contained the complete primary amino acid sequences of their components and thus the complete repertoire of T cell epitopes of the single allergens. The presence of B cell epitopes was investigated with antibodies of predefined specificity for the individual components and by immunological competition experiments. Figure 4 shows that the "giant" is recognized by antisera raised against rPhl p 1, rPhl p 2, rPhl p 5 and rPhl p 6, respectively (Figure 4, row A: 1-4). The correct expression of the C-terminal hexahistidine tag was demonstrated by the reactivity of a mouse monoclonal anti-His tag antibody which specifically recognized the "Giant" (Figure 4: row A, 5) but not rPhl p 2 containing no hexahistidine tag (Figure 4: row B, 5). Recombinant Phl p 2 (row B: 1, 3, 4) and rPhl p 5 (row B: 2) (negative controls) did not react with the antisera (Figure 4: row B).

Next we investigated whether the hybrid allergens can be used to block the binding of grass pollen allergic patients (n=20) IgE antibodies to the individual components (Tables 1, 2). Table 1 shows that the rPhl p 2/rPhl p 6 as well as the rPhl p 6/rPhl p 2 hybrid inhibited IgE binding to both of the components (rPhl p 2, rPhl p 6) in a comparable manner as the single recombinant allergens (approximately 80% inhibition of IgE binding). Likewise, we found that the rPhl p 5/rPhl p 1 hybrid inhibited IgE binding of the 20 patients to both, rPhl p 1 and rPhl p 5, as efficient as the single components (average inhibition 85%) (Table 2). No inhibition was observed when sera were preadsorbed with the rPhl p 5/rPhl p 1 hybrid and reacted with an unrelated allergen (major birch pollen allergen, Bet v 1) whereas rBet v 1 caused almost complete autoinhibition (83% inhibition) (Table 2).

#### Experimental protocol:

#### ELISA competition experiments:

Sera from 20 patients were diluted 1:10 in Tris-buffered saline (TBS) for the inhibition with rPhl p 5/rPhl p 1 and 1:5 for inhibition with rPhl p 2/rPhl p 6 and rPhl p 6/rPhl p 2. The sera were preadsorbed overnight at 4°C either recombinant Phl p 1, Phl p 5, Phl p 5/Phl p 1, Phl p 2, Phl p 6, Phl p 2/Phl p 6, Phl p 6/Phl p 2, Bet v 1, or BSA (10 µg/ml). ELISA plates were coated with recombinant Phl p 1, Phl p 2, Phl p 5, Phl p 6, and Bet v 1 (5 µg/ml) at 4°C overnight. The ELISA was performed as described (Niederberger et al. (1998) J Allergy Clin Immunol 101: 258-264).

#### Dot blots:

Aliquots of approximately 1 µg of purified, recombinant Phl p 6/Phl p 2/Phl p 5/Phl p 1, and rPhl p 5 and rPhl p 2 as negative controls, were dotted on nitrocellulose membranes. Filters were blocked with TBS, 1% BSA, 0.05% Tween-20 for 1 hour and incubated with rabbit anti-Phl p 1, anti-Phl p 5 (diluted 1:5000), anti-Phl p 2 (diluted 1:500), anti-Phl p 6 (diluted 1:2000) sera, and a mouse monoclonal anti-Hexahistidin Ab (Dianova, Germany) diluted 1:1000 in TBS, 1% BSA, 0.05% Tween-20. Filters were washed 3x with TBS, 0.05% Tween-20, incubated with a secondary alkaline phosphatase-coupled goat anti-rabbit IgG antibody (Jackson) and for the His-tag detection with a alkaline phosphatase-coupled anti-mouse IgG



antibody (PharMingen) diluted 1:1000 in TBS, 1%BSA, 0.05% Tween-20 for 1 hour, and washed 3x with TBS, 0.05% Tween-20. Colour reaction was started by adding NBT/BCIP (300µg/ml) and stopped by the addition of water.

Patient	rPhl p 2	rPhl p 2/ rPhl p 6	rPhl p 6/ rPhl p 2	rPhl p 6	r Phl p 2/ rPhl p 6	rPhl p 6/ rPhl p 2
1	69	72	78	72	72	73
2	72	78	81	81	80	81
3	97	95	98	93	91	97
4	93	93	94	80	80	80
5	79	79	78	69	76	75
6	44	44	45	80	82	82
7	97	97	97	87	97	96
8	92	93	92	84	93	93
9	84	85	85	64	68	68
10	53	56	56	45	53	52
11	83	84	82	78	93	91
12	67	64	65	70	72	71
13	90	89	89	86	91	91
14	94	94	93	85	89	89
15	81	81	81	86	89	95
16	97	96	97	95	96	96
17	97	97	94	95	97	97
18	90	90	91	43	53	52
19	76	75	76	68	83	82
20	97	97	97	73	89	89
mean	83	83	84	77	82	83

rPhl p 2 coated

rPhl p 6 coated

Table 1.

Patient	rPhl p 1	rPhl p 5/ rPhl p 1	rPhl p 5	rPhl p 5/ rPhl p 1	rBet v 1	rPhl p 5/ rPhl p 1
1	77	81	85	83	74	0
2	90	89	94	92	96	18
3	96	92	96	92	67	0
4	92	92	95	90	-	-
5	82	84	87	87	-	-
6	92	93	93	93	75	9
7	88	90	90	91	97	0
8	90	91	93	92	88	2
9	50	52	57	57	59	4
10	89	89	93	90	58	5
11	88	95	89	94	-	-
12	95	95	95	96	97	0
13	94	96	94	95	97	0
14	95	96	95	95	98	0
15	78	78	81	79	92	0
16	94	95	95	94	94	0
17	46	58	66	65	-	-
18	77	78	86	85	63	42
19	91	91	94	93	-	-
20	95	93	97	95	96	32
<b>mean</b>	<b>85</b>	<b>86</b>	<b>89</b>	<b>88</b>	<b>83</b>	<b>7</b>

---

 rPhl p 1 coated

---

 rPhl p 5 coated

---

 rBet v 1 coated

23. Dez. 2000

SEQUENCE LISTING

<110> SHAN-Beteiligungsgesellschaft m.b.H.

<120> Allergy vaccines containing hybrid polypeptides

<130> Allergy vaccines containing hybrid polypeptides

<140>

<141>

<150>

<151>

<160> 18

<170> PADAT Sequenzmodul, Version 1.0

<210> 1  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 1

ggaattcata tggccgatct cggttac

27

<210> 2  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 2

cggggtaccg actttgtagc caccagt

27

<210> 3  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 3

cggggtacca tgatccccaa gggtccc

27

<210> 4  
<211> 47  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 4

cgggatcctc agtgggtggtg gtgggtggtgc ttggactcgt agctggt

47

<210> 5  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 5

ggaattcata tgggtgccgaa ggtgacg

27

<210> 6  
<211> 39  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 6

cgtggccttc cccataagct tctcttctgg cgcgtaggt

39

<210> 7  
<211> 24  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 7

aagcttatgg ggaaggccac gacc

24

<210> 8  
<211> 46  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 8

cgggatccta gtggtggtgg tgggtggtgcg cgccgggctt gacagc

46

<210> 9  
<211> 27  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 9

ggaattcata tgggtgccgaa ggtgacg

27

<210> 10  
<211> 46  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 10

cgggatccta gtggtggtgg tgggtggtgcg cgccgggctt gacagc

46

<210> 11  
<211> 30  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<400> 11

ggaattcata tggggaaggc cacgaccgag

30

<210> 12  
<211> 39  
<212> DNA  
<213> artificial sequence

<220>  
<223> primer

<223> primer

<400> 16

cgggatccta gtggtggtgg tgggtggtgct cttctggcgc gtaggt

46

<210> 17

<211> 30

<212> DNA

<213> artificial sequence

<220>

<223> primer

<400> 17

ggaattcata tggggaaggc cacgaccgag

30

<210> 18

<211> 31

<212> DNA

<213> artificial sequence

<220>

<223> primer

<400> 18

gggatttcca tatgctcttc tggcgcgtag g

31



<400> 12

caccttcggc accataagct tcgcgccggg cttgacagc

39

<210> 13

<211> 24

<212> DNA

<213> artificial sequence

<220>

<223> primer

<400> 13

aagcttatgg tgccgaaggt gacg

24

<210> 14

<211> 46

<212> DNA

<213> artificial sequence

<220>

<223> primer

<400> 14

cgggataccta gtggtggtgg tggtggtgct cttctggcgc gtaggt

46

<210> 15

<211> 30

<212> DNA

<213> artificial sequence

<220>

<223> primer

<400> 15

ggaattcata tggggaaggc cacgaccgag

30

<210> 16

<211> 46

<212> DNA

<213> artificial sequence

<220>

EPO-Munich  
52  
28. Dez. 2000

SHAN Beteiligungsgesellschaft m.b.H.

Claims:

1. A hybrid polypeptide comprising at least two different allergenic proteins or fragments thereof wherein each fragment consists of at least eight consecutive amino acids of the respective allergenic protein.
2. A hybrid polypeptide according to claim 1 wherein the hybrid polypeptide comprises at least one complete allergenic protein.
3. A hybrid polypeptide according to claim 2 wherein the hybrid polypeptide comprises at least two complete allergenic proteins.
4. A hybrid polypeptide according to claim 1 wherein the hybrid polypeptide comprises at least one fragment of an allergenic protein which fragment has a substantially reduced allergenic activity compared with the allergenic protein from which it is derived.
5. A hybrid polypeptide according to claim 4 wherein the hybrid polypeptide comprises fragments of at least two different allergenic proteins all of which fragments have a substantially reduced allergenic activity compared with the respective allergenic proteins from which they are derived.
6. A hybrid polypeptide according to any of claims 1-5 characterized in that it comprises at least three different allergenic proteins or fragments thereof.
7. A polynucleotide encoding a polypeptide according to any of claims 1-6.
8. The use of a polynucleotide according to claim 7 for the preparation of a DNA vaccine.
9. A cell transfected or transformed with a polynucleotide comprising a polynucleotide according to claim 7.

10. A pharmaceutical composition containing a polypeptide according to any of claims 1-6, a polynucleotide according to claim 7 or a cell according to claim 9.

11. A pharmaceutical composition according to claim 10 which is a vaccine composition.

12. A pharmaceutical composition according to claim 10 or 11 further containing an adjuvant.

13. A method for the preparation of a hybrid polypeptide according to any of claims 1-6 comprising the following steps:

- a) providing a polynucleotide encoding the hybrid polypeptide;
- b) introducing said polynucleotide into a host cell; and
- c) culturing the host cell obtained in step b) under conditions such that the hybrid polypeptide is expressed; and
- d) recovering the expression product from the cell.

14. A method according to claim 13 wherein the polynucleotide encoding the hybrid polypeptide is prepared using PCR technology.

15. A method for the preparation of a hybrid polypeptide according to any of claims 1-6 characterized in that the polypeptide is prepared by chemical synthesis.

16. The use of a hybrid polypeptide according to any of claims 1-6, or of a polynucleotide according to claim 7, or of a cell according to claim 9 for the preparation of a medicament for the treatment of an allergic disorder.

17. The use of a hybrid polypeptide according to any of claims 1-6, or of a polynucleotide according to claim 7 or of a cell according to claim 9 for the preparation of a medicament for prophylactic vaccination or tolerance induction.

18. The use of a hybrid polypeptide according to any of claims 1-6 for the detection of antibodies against an allergenic protein in a sample.

19. The use according to claim 18 wherein the detection is performed by in vitro antibody tests, tests using cells and in vivo tests.

EPO - Munich  
52

28. Dez. 2000

SHAN Beteiligungsgesellschaft m.b.H.

Abstract

Hybrid polypeptides comprising at least two different allergenic proteins or fragments thereof wherein each fragment consists of at least eight consecutive amino acids of the respective allergenic protein are disclosed. The hybrid polypeptides and polynucleotides coding therefor can be used as pharmaceutical compositions, in particular as vaccines.




Figure 1.





Figure 2.

Figure 3.








Figure 4.